

# **Importation into the United States From South Korea of Fresh Cucurbit Fruits for Consumption**

**Qualitative, Pathway-Initiated Pest Risk Assessment**

**October 1996**


**Agency Contact:**

**Biological Assessment and Taxonomic Support  
Plant Protection and Quarantine  
Animal and Plant Health Inspection Service  
U.S. Department of Agriculture  
4700 River Road, Unit 133  
Riverdale, MD 20737-1236**



## Table of Contents

<b>A. Introduction</b> .....	1
<b>B. Risk Assessment Methods</b> .....	1
<b>1. Initiating Event: Proposed Action</b> .....	1
<b>2. Assessment of Weediness Potential of:</b> <i>Citrullus vulgaris</i> .....	3
: <i>Cucumis sativus</i> .....	4
: <i>Cucumis melo</i> .....	5
: <i>Cucurbita maxima</i> .....	6
<b>3. Previous Risk Assessments, Current Status and Pest Interceptions</b> ...	7
<b>4. Pest List: Pests Associated with cucurbit in Korea</b> .....	8
<b>5. List of Quarantine Pests</b> .....	12
<b>6. Quarantine Pests Likely to Follow Pathway (Quarantine Pests Selected         for Further Analysis</b> .....	13
<b>7. Economic Importance: Consequences of Introduction</b> .....	13
<b>8. Likelihood of Introduction</b> .....	14
<b>9. Conclusion: Pest Risk Potential and Phytosanitary Measures</b> .....	15
<b>C. References</b> .....	15



## A. Introduction

This pest risk assessment was prepared by the Animal and Plant Health Inspection Service (APHIS) of the U.S. Department of Agriculture (USDA) to examine plant pest risks associated with the importation into the United States of fresh cucurbit fruits, (*Citrullus vulgaris*, watermelon; *Cucurbita maxima*, squash; *Cucumis sativus*, cucumber; and *Cucumis melo*, oriental melon) grown in South Korea (Suh, 1991; Lee, 1992). Hereafter in this document the use of the term cucurbit(s) will mean the above named species. This is a qualitative pest risk assessment, that is, estimates of risk are expressed in qualitative terms such as high or low as opposed to numerical terms such as probabilities or frequencies.

International plant protection organizations (*e.g.*, North American Plant Protection Organization (NAPPO), International Plant Protection Convention (IPPC) of the United Nations Food and Agriculture Organization (FAO)) provide guidance for conducting pest risk analyses. The methods we used to initiate, conduct, and report this plant pest risk assessment are consistent with guidelines provided by NAPPO, IPPC and FAO. Our use of biological and phytosanitary terms (*e.g.*, introduction, quarantine pest) conforms with the *NAPPO Compendium of Phytosanitary Terms* (NAPPO 1996) and the *Definitions and Abbreviations* (Introduction Section) in *International Standards for Phytosanitary Measures, Section 1—Import Regulations: Guidelines for Pest Risk Analysis* (FAO 1996).

Pest risk assessment is one component of an overall pest risk analysis. The *Guidelines for Pest Risk Analysis* provided by FAO (1996) describe three stages in pest risk analysis. This document satisfies the requirements of FAO Stages 1 (initiation) and 2 (risk assessment).

The Food and Agriculture Organization (FAO, 1996) defines "pest risk assessment" as "Determination of whether a pest is a quarantine pest and evaluation of its introduction potential". "Quarantine pest" is defined as "A pest of potential economic importance to the area endangered thereby and not yet present there, or present but not widely distributed and being officially controlled" (FAO, 1996; NAPPO, 1996). Thus, pest risk assessments should consider both the likelihood and consequences of introduction of quarantine pests. Both issues are addressed in this qualitative pest risk assessment.

This document presents the findings of our qualitative plant pest risk assessment. We have not described in detail our assessment methods or the criteria we used to rate the various risk elements. Details of our methodology and rating criteria can be found in our "template" document: *Pathway-Initiated Pest Risk Assessment: Guidelines for Qualitative Assessments, version 4.0* (USDA, 1995); to obtain a copy of our template, contact the individual named on the front of this risk assessment.

## B. Risk Assessment

### 1. Initiating Event: Proposed Action

This pest risk assessment is commodity-based, and therefore "pathway-initiated"; we initiated the assessment in response to the request for USDA authorization to allow imports of a

particular commodity presenting a potential plant pest risk. In this case, the importation of certain fresh cucurbit fruits grown in South Korea into the U.S. is a potential pathway for introduction of plant pests. Quarantine 56 (7 CFR §319.56) provides a general regulatory authority for importation of fruits and vegetables.

The cucurbits belong to the Cucurbitaceae family which consists of about 90 genera and about 700 species existing in the tropical regions with some extending into temperate zones. Both annual and perennial species exist within these rapid-growing frost-tender herbs grown for the edible fruits and for ornaments. There are about 25 species in the genus *Cucurbita* [all supposed to be American (Bailey, 1949)], four recognized species of *Citrullus* one of which is widely grown for its edible fruit, some 30 species of *Cucumis* a few of which are grown for the edible fruits.

## 2. Assessment of Weediness Potential of Korean Cucurbits

Tables 1, 1a, 1b, and 1c shows the results of our weediness screening for *Citrullus vulgaris*, *Cucumis sativus*, *Cucumis melo*, and *Cucurbita maxima*. These findings did not require us to initiate a pest-initiated pest risk assessment for any of these species.

**Table 1: Process for Determining Weediness Potential of Commodity**

**Commodity:** *Citrullus vulgaris* = *C. lanatus* var *lanatus* (watermelon)

**Phase 1:** *Citrullus vulgaris* is grown commercially in the U.S and seeds are sold throughout the U.S. for private gardens.

**Phase 2:** Is the species listed in:

Yes *Geographical Atlas of World Weeds* (Holm, 1979)

NO *World's Worst Weeds* (Holm, 1977)

NO *Report of the Technical Committee to Evaluate Noxious Weeds; Exotic Weeds for Federal Noxious Weed Act* (Gunn & Ritchie, 1982)

NO *Economically Important Foreign Weeds* (Reed, 1977)

Yes Weed Science Society of America list (WSSA, 1989)

Yes Is there any literature reference indicating weediness (e.g., *AGRICOLA*, *CAB*, *Biological Abstracts*, *AGRIS*; search on "species name" combined with "weed").

**Phase 3: Conclusion:**

The *Geographical Atlas of World Weeds* includes *Citrullus lanatus* var. *lanatus* as a common weed in Taiwan and a weed of unknown importance in Australia. Weeds included on the Weed Science Society of America list are considered "of current or potential importance in the United States".

Wild watermelon emergence and control. Smith, D. T.; Cooley, A. W., Texas Agric. Exp. Stn., College Stn., 77843, USA. *Weed Science* vol. 21 (6): p.570-573. Publication Year: 1973

Distribution, growth and competitiveness of wild watermelon as a weed. Smith, D. T.; Cooley, A. W.; Hemphill, T. E. Texas A&M Univ., College Station, USA. Miscellaneous Publication, Texas Agricultural Experiment Station (No. MP-1056): 8 pp. Publication Year: 1972

However, this plant does not meet the definition of a Federal Noxious Weed because it is not "new to, or not widely prevalent in the United States."

## Table 1a: Process for Assessing Weediness Potential of Plant Species

**Species:** *Cucumis sativus* L. (cucumber)

**Phase 1:** *Cucumis sativus* is grown commercially in the U.S. and seeds are sold throughout the U.S. for private gardens.

**Phase 2:** Is the species listed in:

Yes *Geographical Atlas of World Weeds* (Holm, 1979)

NO *World's Worst Weeds* (Holm, 1977)

NO *Report of the Technical Committee to Evaluate Noxious Weeds: Exotic Weeds for Federal Noxious Weed Act* (Gunn & Ritchie, 1982)

NO *Economically Important Foreign Weeds* (Reed, 1977)

NO Weed Science Society of America List (WSSA, 1989)

NO Is there any literature reference indicating weediness (e.g., *AGRICOLA*, *CAB*, *Biological Abstracts*, *AGRIS*; search on "species name" combined with "weed").

**Phase 3: Conclusion:**

The *Geographical Atlas of World Weeds* includes *Cucumis sativus* as a common weed in Australia and a weed of unknown importance in Cambodia and India. This plant does not meet the definition of a Federal Noxious Weed because it is not "new to, or not widely prevalent in the United States."

**Table 1b: Process for Assessing Weediness Potential of Plant Species**

**Species:** *Cucumis melo* L. (melon)

**Phase 1:** *Cucumis melo* is grown commercially within the U.S. and seeds are sold throughout the U.S. for private gardens.

**Phase 2:** Is the species listed in:

Yes *Geographical Atlas of World Weeds* (Holm, 1979)

NO *World's Worst Weeds* (Holm, 1977)

NO *Report of the Technical Committee to Evaluate Noxious Weeds: Exotic Weeds for Federal Noxious Weed Act* (Gunn & Ritchie, 1982)

NO *Economically Important Foreign Weeds* (Reed, 1977)

NO Weed Science Society of America List (WSSA, 1989)

NO Is there any literature reference indicating weediness (*e.g.*, *AGRICOLA*, *CAB*, *Biological Abstracts*, *AGRIS*; search on "species name" combined with "weed").

**Phase 3: Conclusion:**

The *Geographical Atlas of World Weeds* includes *Cucumis melo* as a principal weed in Columbia and a weed of unknown importance in Ghana, West Polynesia and Sudan. The Weed Science Society of America includes one variety, *C. melo* var. *dudaim* Naud. (smallmelon).

This species does not meet the definition of a Federal Noxious Weed because it is not "new to, or not widely prevalent in the United States."

**Table 1c: Process for Assessing Weediness Potential of Plant Species**

**Species:** *Cucurbita maxima* Duchesne

**Phase 1:** *Cucurbita maxima* is grown commercially in the U.S. and seeds are sold throughout the U.S. for private gardens.

**Phase 2:** Is the species listed in:

NO *Geographical Atlas of World Weeds* (Holm, 1979)

NO *World's Worst Weeds* (Holm, 1977)

NO *Report of the Technical Committee to Evaluate Noxious Weeds: Exotic Weeds for Federal Noxious Weed Act* (Gunn & Ritchie, 1982)

NO *Economically Important Foreign Weeds* (Reed, 1977)

NO Weed Science Society of America List (WSSA, 1989)

NO Is there any literature reference indicating weediness (e.g., *AGRICOLA*, *CAB*, *Biological Abstracts*, *AGRIS*; search on "species name" combined with "weed").

**Phase 3: Conclusion:**

Species within the genus *Cucurbita* are considered native to the U.S. and *C. maxima* does not meet the definition of a Federal Noxious Weed because it is not "new to, or not widely prevalent in the United States."

### 3. Previous Risk Assessments, Current Status and Pest Interceptions

#### 3a. Decision history for cucurbit fruits from East Asia:

- 1992 - Korea: Cucurbits - Entry denied into U.S., no treatment available for Bactrocera depressus (Shiraki).
- 1986 - Japan: Cucumis melo - Entry denied into U.S., no treatment available for Dacus depressus.
- 1971 - Japan & Korea: Cucurbits approved entry into Guam subject to inspection.
- 1970 - Japan: Cucumis sativus - Entry permitted into Guam subject to inspection.
- 1970 - Korea: Cucumis sativus - Entry permitted into Guam subject to inspection.
- 1969 - Japan: Cucumis sativus - Entry permitted into Hawaii of cucumbers subject to inspection and certification as hothouse grown on Honshu Island or north thereof.
- 1968 - Japan: Cucumis melo - Entry permitted into Hawaii of muskmelons subject to inspection and certification as hothouse grown on Honshu Island or north thereof.
- 1967 - Japan: Cucurbita maxima - Entry permitted into Guam subject to inspection.
- 1967 - Korea: Cucurbita maxima - Entry permitted into Guam subject to inspection.
- 1961 - Japan: Cucurbits spp. - Entry denied into Alaska: multiple pests.
- 1961 - Japan: Cucumis sativus - Entry denied into Alaska: multiple pests.
- 1924 - China: Cucumbers - Entry denied.
- 1924 - Japan: Cucumbers - Entry denied.
- 1924 - Japan: Cucurbita maxima - Entry denied.
- 1923 - China: Squash & pumpkin - Entry denied.

3b. Interceptions from 1985 - 1995:

Table 2: Pest Interceptions of cucurbit fruits from Korea		
Pest	Where Found	Country of Origin
<b>Pathogens</b>		
<i>Cladosporium sp.</i>	baggage	Korea
<i>Phomopsis sp.</i>	baggage	Korea
<b>Arthropods</b>		
Pyraustinae, species of.	baggage	Korea

4. Pest List: Pests Associated with certain cucurbits in Korea

Table 3: Pest List - Korean cucurbits fruits				
Scientific Name and Classification	Distribution <sup>1</sup>	Codes <sup>2</sup>	Hosts <sup>3</sup>	References
<b>INSECTS</b>				
<i>Adoxophyes privatana</i> Walker (Lepidoptera: Tortricidae)	KS	a	3	Anon, 1972; Hill, 1987
<i>Agrotis segetum</i> (Denis & Schiffermuller) (Lepidoptera: Noctuidae)	KS	a	3	Anon, 1957; Anon, 1972
<i>Anomala cuprea</i> Hope (Coleoptera: Scarabaeidae)	KS	a	3	Anon, 1972
<i>Aphis gossypii</i> Glover (Hemiptera: Aphidae)	KS,US	c,e,f,y	1,3,5	Anon, 1972; Blackman & Eastop, 1985
<i>Atrachya menetriesi</i> Faldermann (Coleoptera: Chrysomelidae)	KS	a	3	Anon, 1972
<i>Aulacophora femoralis</i> Mostschulsky (Coleoptera: Chrysomelidae)	KS	a	1,3,5	Anon, 1972
<i>Bactrocera depressa</i> Shiraki (Diptera: Tephritidae)	KS	z <sub>1</sub>	2,3,4,7	Kwon, 1985; Han et al, 1994; Kim, 1994
<i>Bourletiella hortensis</i> Fitch (Collembola: Smythuridae)	KS,US	c,f	1,3,5	Anon, 1972

<i>Chrysodeixis eriosoma</i> (Doubleday) (Lepidoptera: Noctuidae)	KS	a	1,2,6	Stout, 1982; Roberts, 1979
<i>Diaphania indica</i> (Saunders) (Lepidoptera: Pyralidae)	KS,US(FL)	h,z <sub>1</sub>	1,2,3,5,6	Anon, 1972; Whittle & Ferguson, 1987
<i>Frankliniella intonsa</i> (Trybom) (Thysanoptera: Thripidae)	KS	e	2,4,6	Woo et al., 1991
<i>Haplothrips chinensis</i> Priesner (Thysanoptera: Thripidae)	KS	a	2,4,6	Woo et al., 1991
<i>Helicoverpa armigera</i> Hubner (Lepidoptera: Noctuidae)	KS	a	3	Avidov & Harpaz, 1969; Anon, 1970's; EPPO/CABI, 1992
<i>Hylemya platura</i> Meigen (Diptera: Anthomyiidae)	KS,US	c,f	1,3,5	Anon, 1972
<i>Loxoblemmus doenitzii</i> Stein (Orthoptera: Gryllidae)	KS	a	1,3,5	Anon, 1972
<i>Lygus lucorum</i> Meyer-Dur (Hemiptera: Miridae)	KS	a	1,3,5	Anon, 1972; Metcalf & Metcalf, 1993
<i>Lygus spinolaj</i> Meyer-Dur (Hemiptera: Miridae)	KS	a	1,3,5	Anon, 1972; Metcalf & Metcalf, 1993
<i>Mamestra brassicae</i> (Linnaeus) (Lepidoptera: Noctuidae)	KS	a	3	Anon, 1972; Ford, 1988
<i>Monolepta dichroa</i> Harold (Coleoptera: Chrysomelidae)	KS	a	1,3,5	Anon, 1972
<i>Myzus persicae</i> Sulzer (Hemiptera: Aphidae)	KS,US	c,e,f,y	1,3,5	Anon, 1972; Blackman & Eastop, 1985
<i>Ostrinia furnacalis</i> (Guenee) (Lepidoptera: Pyralidae)	KS,US(GU)	h,z <sub>1</sub>	7	Nafus & Schreiner, 1991
<i>Phytomyza horticola</i> Goureau (Diptera: Agromyzidae)	KS	a	2,3,6	Spencer, 1973
<i>Spodoptera litura</i> (Fabricius) (Lepidoptera: Noctuidae)	KS	a	6	Stout, 1982; Wallenamier, 1982
<i>Teleogryllus emma</i> Ohmachi & Matsuura (Orthoptera: Gryllidae)	KS	a	1,3,5	Anon, 1986
<i>Thrips setosus</i> Moulton (Thysanoptera: Thripidae)	KS	a,y	6	Woo et al., 1991; EPPO/CABI, 1992
<i>Thrips tabaci</i> Lindeman (Thysanoptera: Thripidae)	KS,US	c,f,y	1,3,5	Anon, 1972; Metcalf & Metcalf, 1993
<b>ACARINA</b>				
<i>Tetranychus urticae</i> Kock (Tetranychidea: Tetranychidae)	KS,US	c,f	1,3,5	Anon, 1972; Metcalf & Metcalf, 1993
<i>Tetranychus kanzawai</i> Kishida (Tetranychidea: Tetranychidae)	KS	a	5,7	Hong, 1994; Kim, 1994

BACTERIA				
<i>Agrobacterium tumefaciens</i> (Smith & Townsend) Conn.	KS,US	a,c	2,4,7	Bradbury, 1986
<i>Erwinia tracheiphila</i> (Smith) Bergey et al.	KS,US	a,c	4,6,7	Anon, 1986; Bradbury, 1986
<i>Pseudomonas syringae</i> pv. <i>lachrymans</i> (Smith & Bryan) Young, Dye & Wilkie	KS,US	c,z <sub>e</sub>	2,6,7	Anon, 1986; Bradbury, 1986
PATHOGENS				
<i>Alternaria cucumerina</i> (Ellis & Ever.) J.A. Elliott (Fungi Imperfecti: Hyphomycetes)	KS,US	a,c,f	6	Anon, 1986; Farr et al, 1986; Ellis & Holliday, 1970
<i>Athelia rolfsii</i> (Curzi) Tu & Kimbrough (Basidiomycetes: Aphyllophorales)	KS,US	a,c,f	1,6,7	Anon, 1972; Farr et al, 1989
<i>Botrytis cinerea</i> Pers.:Fr. (Fungi Imperfecti: Hyphomycetes)	KS,US	a,c,f	3,5,6	Anon, 1986; Farr et al, 1989
<i>Cercospora citrullina</i> Cooke (Fungi Imperfecti: Hyphomycetes)	KS,US	a,c,f	1-7	Anon, 1986
<i>Cladosporium cucumerinum</i> Ellis & Arthur (Fungi Imperfecti: Hyphomycetes)	KS,US	c,f,z <sup>a</sup>	2,6,7	Anon, 1986; Farr et al, 1989; Jarvis, 1992
<i>Colletotrichum orbiculare</i> (Berk. & Mont.) Arx (Fungi Imperfecti: Coelomycetes)	KS,US	c,f,z <sub>e</sub>	1,2,3,5,6,7	Anon, 1986; Farr et al, 1989
<i>Didymella bryoniae</i> (Auersw.) Rehm (Loculoascomycetes: Dothideales)	KS,US	c,f,z <sub>e</sub>	1,2,3,5,7	Anon, 1986; Farr et al, 1989
<i>Erysiphe cichoracearum</i> DC. (Pyrenomycetes: Erysiphales)	KS,US	a,c,f	1,2,3,5	Anon, 1986; Farr et al, 1989
<i>Fusarium oxysporum</i> Schlechtend.:Fr. f. sp. <i>cucumerinum</i> J. H. Owen (Fungi Imperfecti: Hyphomycetes)	KS,US	a,c,f	5,6	Anon, 1986; Farr et al, 1989
<i>Fusarium oxysporum</i> Schlechtend.:Fr. f. sp. <i>melonis</i> Snyder & Hansen (Fungi Imperfecti: Hyphomycetes)	KS,US	a,c,f	5,7	Anon, 1986; Farr et al, 1989
<i>Fusarium oxysporum</i> Schlechtend.:Fr. f. sp. <i>niveum</i> (Smith) Snyder & Hansen (Fungi Imperfecti: Hyphomycetes)	KS,US	a,c,f	1,2,3,5	Anon, 1986; Farr et al, 1989
<i>Phytophthora melonis</i> Katsura (Oomycetes: Peronosporales)	KS	a	6	Anon, 1986
<i>Phytophthora nicotianae</i> Breda de Hann var. <i>parasitica</i> (Dastur) Waterhouse (Oomycetes: Peronosporales)	KS,US	c,f,z <sub>e</sub>	1,3,6,7	Anon, 1986; Farr et al, 1989; Waterhouse & Waterhouse, 1964
<i>Pseudoperonospora cubensis</i> (Berk. & M.A. Curtis) Rostovzev (Oomycetes: Peronosporales)	KS,US	a,c,f	1,3,5,6,7	Anon, 1986; Farr et al, 1989; Jarvis, 1992; Palti, 1975
<i>Pythium debaryanum</i> Hesse (Oomycetes: Peronosporales)	KS,US	a,c,f	1,3,5,6,7	Anon, 1986; Farr et al, 1989
<i>Rhizoctonia solani</i> Kuhn (Fungi Imperfecti: Agonomycetes)	KS,US	a,c,f	6,7	Anon, 1986; Farr et al, 1989; Mordue, 1974

<i>Sphaerotheca fuliginea</i> (Schlechtend.:Fr.) Pollacci (Pyrenomycetes: Erysiphales)	KS,US	a,c,f	1,2,3,5,6,7	Anon, 1986; Farr et al., 1989; Jarvis, 1992
<b>VIRUSES</b>				
Bean yellow mosaic virus	KS,US	a,c,f	2	Bos, 1970; Ryu et al., 1986; Lee et al., 1983
Cucumber green mottle mosaic virus	KS	z <sup>1</sup>	1,3,5,6,7	Hollings et al., 1975; Lee et al., 1990
Cucumber mosaic virus	KS,US	c,f	2,3,6,7	Anon, 1972
Tobacco ringspot virus	KS,US	c,f	3,6	Stace-Smith, 1985; Cho et al., 1976
Watermelon mosaic 2 potyvirus	KS,US	a,c,f	2,4,6,7	Brunt et al., 1990; Purcifull et al., 1984
<b>NEMATODES</b>				
<i>Meloidogyne arenaria</i> (Neal) Chitwood	KS,US	a,c	1,3,5	Anon, 1972; Anon, 1985
<i>Meloidogyne incognita</i> (Kofoid & White) Chitwood	KS,US	a,c	1,3,5	Anon, 1972; Anon, 1985

Distribution legend: KS = Korea; US = United States; FL = Florida; GU = Guam

Codes:

- a = Pest mainly associated with plant part other than commodity
- c = Organism does not meet the geographical and regulatory definition for a quarantine pest.
- e = Although pest attacks commodity, it would not be expected to remain with the commodity during processing
- f = Pest occurs in the U.S. and is not subject to official restrictions and regulations
- h = Pest is present in the U.S. and is listed in the USDA catalogue of intercepted pests as actionable at ports of entry, but, the pest is not currently subject to further official restrictions and regulations
- y = Pest is a vector of plant pathogen(s)
- z<sub>i</sub> = Internal: Pest is known to commonly attack or infect commodity and it would be reasonable to expect the pest may remain with the fruit during processing and shipping
- z<sub>e</sub> = External: Pest is known to infect or infest the commodity and it would be reasonable to expect the pest may remain with the commodity during processing and shipping

Host Code: *Citrullus* sp. = 1; *Citrullus vulgaris* = 2; *Cucurbita* sp. = 3; *Cucurbita maxima* = 4; *Cucumis* sp. = 5; *Cucumis sativus* = 6; *Cucumis melo* = 7

## 5. List of Quarantine Pests

Our list of quarantine pests for commercial shipments of cucurbits from Korea is provided in Table 4. Should any of these pests be intercepted on commercial (or any other) shipments of cucurbits, quarantine action may be taken.

**Table 4: Quarantine Pests: Korean cucurbit fruits for consumption**

<b>Arthropods</b>	<i>Adoxophyes privatana</i> Walker
	<i>Agrotis segetum</i> (Denis & Schiffermuller)
	<i>Anomala cuprea</i> Hope
	<i>Atrachya menetriesi</i> Mostschulsky
	<i>Bactrocera depressa</i> Shiraki
	<i>Chrysodeixis eriosoma</i> (Doubleday)
	<i>Frankliniella intonsa</i> (Trybom)
	<i>Haplothrips chinensis</i> Priesner
	<i>Helicoverpa armigera</i> Hubner
	<i>Loxoblemmus doenitzi</i> Stein
	<i>Lygus lucorum</i> Meyer-Dur
	<i>Mamestra brassicae</i> (Linnaeus)
	<i>Monolepta dichroa</i> Harold
	<i>Ostrinia furnacalis</i> (Guenee)
	<i>Phytomyza horticola</i> Goureau
	<i>Spodoptera litura</i> (Fabricius)
	<i>Teleogryllus emma</i> Ohmachi & Matsuura
<i>Thrips setosus</i> Moulton	
<b>Acarina</b>	<i>Tetranychus kanzawaii</i> Kishida
<b>Pathogen</b>	<i>Phytophthora melonis</i> Katsura
<b>Viruses</b>	Cucumber green mottle mosaic virus

6. **Quarantine Pests Likely to Follow Pathway (i.e., Quarantine Pests Selected for Further Analysis)**

We analyzed in detail only those quarantine pests that can reasonable be expected to follow the pathway, i.e., be included in commercial shipments of cucurbit fruits (see USDA, 1995) for selection criteria). Only quarantine pests selected for further analysis are subject to steps 7-9 below.

Table 5: Quarantine Pest Selected for Further Analysis: Korean cucurbits fruits for consumption	
<b>Arthropods</b>	<i>Bactrocera depressa</i> Shiraki <i>Diaphania indica</i> (Saunders) <i>Ostrinia furnacalis</i> (Guenee) <i>Tetranychus kanzawai</i> Kishida
<b>Viruses</b>	Cucumber green mottle mosaic virus

7. **Economic Importance: Consequences of Introduction**

For each quarantine pest selected for further analysis, we consider the consequences of introduction. For qualitative, pathway-initiated pest risk assessments, these risks are estimated by rating each pest with respect to five risk elements. A full description of these elements and rating criteria can be found in USDA (1995). Table 6 shows our risk ratings for these risk elements.

Table 6: Risk Rating: Consequences of Introduction						
Pest	Climate/Host	Host Range	Dispersal	Economic	Environmental	Risk Rating
<i>Bactrocera depressa</i>	high	medium	low	medium	high*	medium
<i>Diaphania indica</i>	high	medium	high	medium	high*	high
<i>Ostrinia furnacalis</i>	high	high	high	medium	low	high
<i>Tetranychus kanzawai</i>	high	high	low	low	low	medium
Cucumber green mottle mosaic virus	medium	low	low	medium	high*	medium

\*This pest is known to attack members of the plant genus *Cucurbita*. In the United States, *Cucurbita okeechobeensis* is a Federally listed endangered species. We believe it would be reasonable to assume that this pest would attack this endangered plant. Because of existing legislation regarding endangered plants, we automatically give this pest a risk rating of "high" for consequences of introduction.

## 8. Likelihood of Introduction

The undesirable outcome being considered is the introduction of a quarantine pest. The two general components of a pest risk assessment are estimates of the consequences and likelihood of introduction. Risk Elements in Table 7 focus on the consequences of introduction. For qualitative pest risk assessments the assessment is as shown below. There are two separate components. First, an estimate is made concerning the amount of commodity likely to be imported; the result is a risk rating (0-2) that applies to the commodity and country in question and is the same for all quarantine pest considered. And second, five biological features concerning the pest and their interactions with the commodity are considered. The resulting risk ratings are specific to each pest.

<b>Table 7: Risk Rating: Likelihood of Introduction</b>						
<b>Pest</b>	<b>Quantity of commodity imported annually</b>	<b>Likelihood survive postharvest treatment</b>	<b>Likelihood survive shipment</b>	<b>Likelihood not detected at port of entry</b>	<b>Likelihood moved to suitable habitat</b>	<b>Likelihood find suitable host</b>
<i>Bactrocera depressa</i>	medium	high	high	high	high	high
<i>Diaphania indica</i>	medium	high	high	medium	high	medium
<i>Ostrinia furnacalis</i>	medium	high	high	medium	high	medium
<i>Tetranychus kanzawai</i>	medium	medium	medium	high	medium	low
Cucumber green mottle mosaic virus	medium	high	high	high	low	low

## 9. Conclusion: Pest Risk Potential and Phytosanitary Measures

Our measure of pest risk potential combines the risk ratings for consequences and likelihood of introduction as described in USDA (1995). Table 8 shows our estimated pest risk potential for the quarantine pests selected for further analysis for importation of Korean cucurbits.

<b>Pest</b>	<b>Pest risk potential</b>
<i>Bactrocera depressa</i>	high
<i>Diaphania indica</i>	high
<i>Ostrinia furnacalis</i>	high
<i>Tetranychus kanzawai</i>	medium
Cucumber green mottle mosaic virus	medium

We recommend specific phytosanitary measures for pest receiving a high PRP risk rating; port-of-entry inspection is not considered sufficient to provide phytosanitary security. APHIS has not yet determined whether risks associated with the importation of the cucurbit fruits listed in this document can be managed adequately. Detailed examination and choice of appropriate sanitary and phytosanitary measures to mitigate pest risk is undertaken as part of the pest risk management phase and is not discussed in this document. Appropriate sanitary and phytosanitary measures to mitigate pest risk will be determined during the pest risk management phase.

## C. References

- Anonymous. 1957. Insects Not Known to Occur in the United States, Number 25. *Agrotis segetum*.
- Anonymous. 1970's. Pathway studies, *Heliothis armigera*. Unpublished USDA, APHIS, PPQ document. p.33.
- Anonymous. 1972. A list of plant diseases, insect pests, and weeds in Korea. Korean Society of Plant Protection. 424 pp
- Anonymous. 1985. Distribution of plant-parasitic nematode species in North America. Society of Nematologists. 205 pp.
- Anonymous. 1986. A List of Plant Diseases, Insects Pests, and Weeds in Korea. 2nd Edition, Korean Society of Plant Protection. 633 pp
- Avidov, Z. & I. Harpaz. 1969. Plant Pests of Israel. Israel Universities Press, Jerusalem, Israel. 548 pp
- Bailey, L. H. 1949. Manual of Cultivated Plants, Most commonly grown in the Continental United States and Canada. 1116 pp

- Blackman, R. L. & V. F. Eastop. 1985. *Aphids on the World's Crops: An Identification Guide*. J. Wiley & Sons, NY, NY. 466 pp.
- Bos, L. 1970. Bean Yellow Mosaic Virus. CMI/AAB descriptions of Plant Viruses No. 40. Commonwealth Mycological Institute/ Association Applied Biologist, Kew, Surrey, England. 4pp.
- Bradbury, J. F. 1986. *Guide to Plant Pathogenic Bacteria*. CAB International Mycological Institute. 332 pp.
- Brunt, A.; K. Crabtree and A. Gibbs. 1990. *Viruses of Tropical Plants*. CAB International, UK. 707 pp.
- Cho, E. K. & B. J. Chung. 1976. Studies on the identification and classification of soybean virus diseases in Korea. I. Preliminary studies on a soybean virus disease. *Korean J. Plant Protection* 15: 61-68.
- Ellis, M. B. & P. Holliday. 1970. CMI Description of Pathogenic Fungi and Bacteria, No. 244. *Alternaria cucumerina*. Commonwealth Mycological Institute, England.
- EPPO/CABI. 1992. *Helicoverpa armigera*. In: Quarantine pests for Europe (Ed. by Smith, I.M.; McNamara, D.G.; Scott, P.R.; Harris, K.M.). CAB International, Wallingford, U.K.
- EPPO/CABI. 1992. Tomato spotted wilt virus. In: Quarantine pests for Europe (Ed. by Smith, I.M.; McNamara, D.G.; Scott, P.R.; Harris, K.M.). CAB International, Wallingford, U.K.
- FAO. 1996. *International Standards for Phytosanitary Measures. Section 1 - Import Regulations: Guidelines for Pest Risk Analysis (Draft Standard)*. Secretariate of the International Plant Protection Convention of the Food and Agriculture Organization of the United Nations. Rome, Italy.
- Farr, D. F., G. F. Bills, G. P. Chamuris and A. Y. Rossman. 1989. *Fungi on plants and plant products in the United States*. American Phytopathological Society, St. Paul, Minnesota. 1252 pp.
- Ford, E. J. 1988. Pathway Studies, *Mamestra brassicae*. Unpublished USDA, APHIS, PPQ document. p. 202-208.
- Gunn, C.R. and C. Ritchie. 1982. 1982 Report of the Technical Committee to Evaluate Noxious Weeds; Exotic Weeds for Federal Noxious Weed Act. (unpublished).
- Han, M.J.; S.H. Lee, S.B. Ahn; J.Y. Choi and K.M. Choi. 1994. Distribution, damage and host plants of pumpkin fruit fly, *Paradacus depressus*. *RDA Journal of Agricultural Science Crop Protection* 36(1). p. 346-350.
- Hill, D. S. 1987. *Agricultural insect pests of temperate regions and their control*. Cambridge University Press. London, England. 659 pp.
- Hollings, M.; Y. Komuro; H. Tochiwara. 1975. Cucumber Green Mottle Mosaic Virus. CMI/AAB Descriptions of Plant Viruses No. 154. Commonwealth Mycological Institute/ Association Applied Biologists, Kew, Surrey, England. 4 pp.
- Holm, L.G., D.L. Plucknett, J.V. Pancho and J.P. Herberger. 1977. *The World's Worst Weeds*. University of Hawaii Press, Honolulu.
- Holm, L.G., J.V. Pancho and J.P. Herberger and D.L. Plucknett. 1979. *A Geographical Atlas of World Weeds*. John Wiley and Sons, New York.
- Hong, I.H. 1994. Official letter to B. G. Lee, Deputy Administrator, PPQ, APHIS, USDA.
- Jarvis, W. R. 1992. *Cucumber diseases*. Agriculture Canada Publication 1684/E. Ottawa, Ont. 49 pp.
- Kim, Y. H. 1994. FAX to E. Imai transmitting information on Cucurbits from Dr. Kwon, Kyongbuk University, Daegu, Korea.

- Kwon, Y. J. 1985. Classification of the fruit fly pests from Korea - *Insecta Koreana* (Series 5) 49-112.
- Lee, B. G. 1992. February letter to Ki Ho Shu, Director, International Quarantine Information Division, Korea.
- Lee, K. W.; B. C. Lee; H. C. Park and Y. S. Lee. 1990. Occurrence of cucumber green mottle mosaic virus disease of watermelon in Korea. *Korean J. Plant Pathology* 6:250-255.
- Lee, S. H., J. S. Kim, and Y. M. Choi. 1983. Bean yellow mosaic virus and cucumber mosaic virus causing mosaic disease on gladiolus in Korea. *Korean Journal of Plant Protection*. 22:3, p. 198-202.
- Metcalf, R. L. & R. A. Metcalf. 1993. *Destructive and Useful Insects, Their Habits and Control*. 5th Edition. p. 14-16.
- Mordue, J. E. M. 1974. CMI Description of Plant Pathogenic Fungi and Bacteria No. 406. *Rhizoctonia solani*. Commonwealth Mycological Institute. England
- Nafus, D. M. & I. H. Schreiner. 1991. Review of the biology & control of the Asian corn borer, *Ostrinia furnacalis* (Lep:Pyralidae). *Tropical Pest Management* 37(1) p. 41-56.
- NAPPO. 1996. NAPPO Compendium of Phytosanitary Terms.. North American Plant Protection Organization (NAPPO). NAPPO Secretariate, Ottawa, Ontario, Canada. 25 pp.
- Palti, J. 1975. CMI Description of Pathogenic Fungi and Bacteria, No. 457. *Pseudoperonospora cubensis*. Commonwealth Mycological Institute, England.
- Purcifull, D. E.; E. Hiebert & J. Edwardson. 1984. CMI Description of Plant Viruses No. 293, Watermelon mosaic virus 2.
- Reed, C.F. 1977. Economically Important Foreign Weeds. Agriculture Handbook No. 498
- Roberts, L. I. N. 1979. Biology of *Chrysodeixis eriosoma* (Lepidoptera: Noctuidae) in New Zealand. *New Zealand Ent.*7(1) p. 52-58.
- Ryu, G. H.; S. H. Lee and Y. J. La. 1986. Identification of bean yellow mosaic virus from *Trifolium repens*. *Korean J. Plant Pathology*. 2: 48-52.
- Spencer, K. A. 1973. *Agromyzidae (Diptera) of Economic Importance*. The Hague, Netherlands. 418 pp.
- Stace-Smith, R. 1985. Tobacco Ringspot Virus. Descriptions of Plant Virus Number 309, Commonwealth Mycological Institute/Association Applied Biologist. Kew, Surrey, Eng. 6 p.
- Stout, O.O. 1982. Plant Quarantine Guidelines within the Pacific Region. p. 656.
- Suh, Ki-Ho 1991. Letter to B. G. Lee, Deputy Administrator, PPQ, APHIS, USDA.
- USDA. 1995. Pathway-Initiated Pest Risk Assessment: Guidelines for Qualitative Assessments, ver 4.0 PPQ, APHIS. 15 pp.
- Wallenmaier, T. 1982. Pathway Studies, *Spodoptera litura*. Unpublished USDA, APHIS, PPQ document. p. 196-197.
- Waterhouse, G. M. & J. M. Waterhouse. 1964. CMI Description of Plant Pathogenic Fungi and Bacteria, No. 35. *Phytophthora nicotianae* var. *parasitica*. Commonwealth Mycological Institute, England.
- Whittle, K. & D. C. Ferguson. 1987. Pest not known to occur in the United States, *Diaphania indica*. APHIS 81-50, USDA. 8 pp.
- Woo, K. S.; O-K. & K-S Cho. 1991. Studies on the distribution, host plants and taxonomy of Korean thrips (Insecta: Thysanoptera). *Seoul Nat'l Univ. J. Agric. Sci.* 16(2) pp. 133-145.
- WSSA, 1989. Composite List of Weeds. Weed Science Society of America.

John W. Lightfield  
Biological Assessment and Taxonomic Support  
Operational Support  
Plant Protection and Quarantine

Reviewers:

G. Cave, Entomologist  
M. Firko, Entomologist\*  
E. Podleckis, Plant Virologist\*  
L. Redmond, Plant Pathologist\*  
S. Redlin, Plant Pathologist\*  
R. Stewart, Entomologist\*